

Extensions of Quandles and Cocycle Knot Invariants

J. Scott Carter
University of South Alabama
Mobile, AL 36688
carter@mathstat.usouthal.edu

Mohamed Elhamdadi
University of South Florida
Tampa, FL 33620
emohamed@math.usf.edu

Marina Appiou Nikiforou
University of South Florida
Tampa, FL 33620
mappiou@chuma.cas.usf.edu

Masahico Saito
University of South Florida
Tampa, FL 33620
saito@math.usf.edu

February 8, 2008

Abstract

Quandle cocycles are constructed from extensions of quandles. The theory is parallel to that of group cohomology and group extensions. An interpretation of quandle cocycle invariants as obstructions to extending knot colorings is given, and is extended to links component-wise.

1 Introduction

A quandle is a set with a self-distributive binary operation (defined below) whose study has been partially motivated from knot theory. A (co)homology theory was defined in [3] for quandles, which is a modification of rack (co)homology defined in [11]. State-sum invariants using quandle cocycles as weights are defined [3] and computed for important families of classical knots and knotted surfaces [4]. Quandle homomorphisms and virtual knots are applied to this homology theory [5]. The invariants were applied to study knots, for example, in detecting non-invertible knotted surfaces [3]. On the other hand, knot diagrams colored by quandles can be used to study quandle homology groups. This view point was developed in [11, 12, 15] for rack homology and homotopy, and generalized to quandle homology in [6].

In [7], constructions of extensions of quandles using cocycles are given, which are similar to extensions of groups using group cocycles [2]. In this paper, we develop methods of constructing cocycles from extensions. This is the opposite direction of [7]. After reviewing the material in Section 2, infinite families of extensions of Alexander quandles are given, and explicit formulas of computing corresponding cocycles are established in Section 3. We show that these families

of extensions are non-trivial. In Section 4, an interpretation of the state-sum cocycle invariants as obstructions to extending colorings by a quandle to colorings by another quandle is given. Generalizations to links are also considered.

2 Quandle and Their Homology Theory

In this section we review necessary material from the papers mentioned in the introduction.

A *quandle*, X , is a set with a binary operation $(a, b) \mapsto a * b$ such that

- (I) For any $a \in X$, $a * a = a$.
- (II) For any $a, b \in X$, there is a unique $c \in X$ such that $a = c * b$.
- (III) For any $a, b, c \in X$, we have $(a * b) * c = (a * c) * (b * c)$.

A *rack* is a set with a binary operation that satisfies (II) and (III).

Racks and quandles have been studied in, for example, [1, 9, 17, 18, 20].

The axioms for a quandle correspond respectively to the Reidemeister moves of type I, II, and III (see [9, 18], for example).

A function $f : X \rightarrow Y$ between quandles or racks is a *homomorphism* if $f(a * b) = f(a) * f(b)$ for any $a, b \in X$.

The following are typical examples of quandles.

- A group $X = G$ with n -fold conjugation as the quandle operation: $a * b = b^{-n}ab^n$.
- Any set X with the operation $x * y = x$ for any $x, y \in X$ is a quandle called the *trivial* quandle. The trivial quandle of n elements is denoted by T_n .
- Let n be a positive integer. For elements $i, j \in \{0, 1, \dots, n-1\}$, define $i * j \equiv 2j - i \pmod{n}$. Then $*$ defines a quandle structure called the *dihedral quandle*, R_n . This set can be identified with the set of reflections of a regular n -gon with conjugation as the quandle operation.
- Any $\Lambda(= \mathbb{Z}[T, T^{-1}])$ -module M is a quandle with $a * b = Ta + (1 - T)b$, $a, b \in M$, called an *Alexander quandle*. Furthermore for a positive integer n , a *mod- n Alexander quandle* $\mathbb{Z}_n[T, T^{-1}]/(h(T))$ is a quandle for a Laurent polynomial $h(T)$. The mod- n Alexander quandle is finite if the coefficients of the highest and lowest degree terms of h are units in $\mathbb{Z}_n$.

Let $C_n^R(X)$ be the free abelian group generated by n -tuples $(x_1, \dots, x_n)$ of elements of a quandle X . Define a homomorphism $\partial_n : C_n^R(X) \rightarrow C_{n-1}^R(X)$ by

$$\begin{aligned} \partial_n(x_1, x_2, \dots, x_n) &= \sum_{i=2}^n (-1)^i [(x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \\ &\quad - (x_1 * x_i, x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)] \end{aligned} \tag{1}$$

for $n \geq 2$ and $\partial_n = 0$ for $n \leq 1$. Then $C_*^R(X) = \{C_n^R(X), \partial_n\}$ is a chain complex.

Let $C_n^D(X)$ be the subset of $C_n^R(X)$ generated by n -tuples $(x_1, \dots, x_n)$ with $x_i = x_{i+1}$ for some $i \in \{1, \dots, n-1\}$ if $n \geq 2$; otherwise let $C_n^D(X) = 0$. If X is a quandle, then $\partial_n(C_n^D(X)) \subset$




Figure 1: Type III move and the quandle identity

$C_{n-1}^D(X)$ and $C_*^D(X) = \{C_n^D(X), \partial_n\}$ is a sub-complex of $C_*^R(X)$. Put $C_n^Q(X) = C_n^R(X)/C_n^D(X)$ and $C_*^Q(X) = \{C_n^Q(X), \partial'_n\}$, where ∂'_n is the induced homomorphism. Henceforth, all boundary maps will be denoted by ∂_n .

For an abelian group G , define the chain and cochain complexes

$$C_*^W(X; G) = C_*^W(X) \otimes G, \quad \partial = \partial \otimes \text{id}; \quad (2)$$

$$C_W^*(X; G) = \text{Hom}(C_*^W(X), G), \quad \delta = \text{Hom}(\partial, \text{id}) \quad (3)$$

in the usual way, where $W = D, R, Q$.

The n th *quandle homology group* and the n th *quandle cohomology group* [3] of a quandle X with coefficient group G are

$$H_n^Q(X; G) = H_n(C_*^Q(X; G)), \quad H_Q^n(X; G) = H^n(C_Q^*(X; G)). \quad (4)$$

Let a classical knot diagram be given. The co-orientation is a family of normal vectors to the knot diagram such that the pair (orientation, co-orientation) matches the given (right-handed, or counterclockwise) orientation of the plane. At a crossing, if the pair of the co-orientation of the over-arc and that of the under-arc matches the (right-hand) orientation of the plane, then the crossing is called *positive*; otherwise it is *negative*. Crossings in Fig. 1 are positive by convention.

A *coloring* of an oriented classical knot diagram is a function $\mathcal{C} : R \rightarrow X$, where X is a fixed quandle and R is the set of over-arcs in the diagram, satisfying the condition depicted in the top of Fig. 1. In the figure, a crossing with over-arc, r , has color $\mathcal{C}(r) = y \in X$. The under-arcs are called r_1 and r_2 from top to bottom; the normal (co-orientation) of the over-arc r points from r_1 to r_2 . Then it is required that $\mathcal{C}(r_1) = x$ and $\mathcal{C}(r_2) = x * y$.

Note that locally the colors do not depend on the orientation of the under-arc. The quandle element $\mathcal{C}(r)$ assigned to an arc r by a coloring $\mathcal{C}$ is called a *color* of the arc. This definition of colorings on knot diagrams has been known, see [9, 13] for example. Henceforth, all the quandles that are used to color diagrams will be finite.

In Fig. 1 bottom, the relation between Reidemeister type III move and a quandle axiom (self-distributivity) is indicated. In particular, the colors of the bottom right segments before and after the move correspond to the self-distributivity.

A (Boltzmann) weight, $B(\tau, \mathcal{C})$, at a crossing τ is defined as follows. Let $\mathcal{C}$ denote a coloring. Let r be the over-arc at τ , and r_1, r_2 be under-arcs such that the normal to r points from r_1 to r_2 . Let $x = \mathcal{C}(r_1)$ and $y = \mathcal{C}(r)$. Pick a quandle 2-cocycle $\phi \in Z^2(X; A)$. Then define $B(\tau, \mathcal{C}) = \phi(x, y)^{\epsilon(\tau)}$, where $\epsilon(\tau) = 1$ or -1 , if the sign of τ is positive or negative, respectively.

The *partition function*, or a *state-sum*, is the expression

$$\sum_{\mathcal{C}} \prod_{\tau} B(\tau, \mathcal{C}).$$

The product is taken over all crossings of the given diagram, and the sum is taken over all possible colorings. The values of the partition function are taken to be in the group ring $\mathbb{Z}[A]$ where A is the coefficient group written multiplicatively. This is proved [3] to be a knot invariant, called the (*quandle*) *cocycle invariant*.

3 Extensions of quandles by 2-cocycles

In [7], for a quandle X , an abelian group A , and a 2-cocycle $\phi \in Z_Q^2(X; A)$, the *abelian extension* $E = E(X, A, \phi)$ was defined as $A \times X$ as a set, with the quandle operation $(a_1, x_1) * (a_2, x_2) = (a_1 \phi(x_1, x_2), x_1 * x_2)$. The following lemma is the converse of the fact proved in [7] that $E(X, A, \phi)$ is a quandle.

Lemma 3.1 *Let X, E be finite quandles, and A be a finite abelian group written multiplicatively. Suppose there exists a bijection $f : E \rightarrow A \times X$ with the following property. There exists a function $\phi : X \times X \rightarrow A$ such that for any $e_i \in E$ ($i = 1, 2$), if $f(e_i) = (a_i, x_i)$, then $f(e_1 * e_2) = (a_1 \phi(x_1, x_2), x_1 * x_2)$. Then $\phi \in Z_Q^2(X; A)$.*

Proof. For any $x \in X$ and $a \in A$, there is $e \in E$ such that $f(e) = (a, x)$, and

$$(a, x) = f(e) = f(e * e) = (a \phi(x, x), x),$$

so that we have $\phi(x, x) = 1$ for any $x \in X$.

By identifying $A \times X$ with E by f , the quandle operation $*$ on $A \times X$ is defined, for any (a_i, x_i) ($i = 1, 2$), by

$$(a_1, x_1) * (a_2, x_2) = (a_1 \phi(x_1, x_2), x_1 * x_2).$$

Since $A \times X$ is a quandle isomorphic to E under this $*$, we have

$$\begin{aligned} & [(a_1, x_1) * (a_2, x_2)] * (a_3, x_3) \\ &= (a_1 \phi(x_1, x_2), x_1 * x_2) * (a_3, x_3) \\ &= (a_1 \phi(x_1, x_2) \phi(x_1 * x_2, x_3), (x_1 * x_2) * x_3), \end{aligned}$$

and

$$\begin{aligned}
& [(a_1, x_1) * (a_3, x_3)] * [(a_2, x_2) * (a_3, x_3)] \\
&= (a_1\phi(x_1, x_3), x_1 * x_3) * (a_2\phi(x_2, x_3), x_2 * x_3) \\
&= (a_1\phi(x_1, x_3)\phi(x_1 * x_3, x_2 * x_3), (x_1 * x_3) * (x_2 * x_3))
\end{aligned}$$

are equal for any (a_i, x_i) ($i = 1, 2, 3$). Hence ϕ satisfies the 2-cocycle condition. ■

This lemma implies that under the same assumption we have $E = E(X, A, \phi)$, where $\phi \in Z_Q^2(X; A)$. Next we identify such examples.

Theorem 3.2 *For any positive integers q and m , $E = \mathbb{Z}_{q^{m+1}}[T, T^{-1}]/(T - 1 + q)$ is an abelian extension $E = E(\mathbb{Z}_{q^m}[T, T^{-1}]/(T - 1 + q), \mathbb{Z}_q, \phi)$ of $X = \mathbb{Z}_{q^m}[T, T^{-1}]/(T - 1 + q)$ for some cocycle $\phi \in Z_Q^2(X; \mathbb{Z}_q)$.*

Proof. Represent elements of $\mathbb{Z}_{q^{m+1}}$ by $\{0, 1, \dots, q^{m+1} - 1\}$ and express them in their q^{m+1} -ary expansion:

$$A = A_m q^m + \dots + A_1 q + A_0 \in \mathbb{Z}_{q^{m+1}},$$

where $0 \leq A_j < q$, $j = 0, \dots, m$. With this convention, A_j 's are uniquely determined integers. Define $f : E \rightarrow \mathbb{Z}_q \times X$ by

$$f(A) = (A_m \pmod{q}, \overline{A} \pmod{q^{m-1}}),$$

where $\overline{A} = \sum_{j=0}^{m-1} A_j q^j$. Then for $A, B \in \mathbb{Z}_{q^{m+1}}$, the quandle operation is computed in $\mathbb{Z}_{q^{m+1}}$ by

$$\begin{aligned}
A * B &= TA + (1 - T)B \\
&= (1 - q)(A_m q^m + \dots + A_1 q + A_0) \\
&\quad + q(B_m q^m + \dots + B_1 q + B_0) \\
&= (A_m - A_{m-1} + B_{m-1})q^m + (A_{m-1} - A_{m-2} + B_{m-2})q^{m-1} \\
&\quad + \dots + (A_1 - A_0 + B_0)q + A_0 \\
&= (A_m - A_{m-1} + B_{m-1})q^m + \sum_{j=0}^{m-1} (A_j - A_{j-1} + B_{j-1})q^j,
\end{aligned}$$

where A_{-1}, B_{-1} are understood to be zeros in the last summation. Define a set-theoretic section $s : \mathbb{Z}_{q^m} \rightarrow \mathbb{Z}_{q^{m+1}}$ by

$$s\left(\sum_{j=0}^{m-1} X_j q^j\right) = 0 \cdot q^m + \sum_{j=0}^{m-1} X_j q^j.$$

For $X, Y \in \mathbb{Z}_{q^m}$ define

$$\phi(X, Y) = [s(X) * s(Y) - s(X * Y)]/q^m \in \mathbb{Z}_q$$

where division by q^m means to consider these elements as integers, divide by q^m and compute the residue class modulo q . Note that $\overline{s(X) * s(Y)} = \overline{s(X * Y)}$. Hence, $s(X) * s(Y) - s(X * Y)$ is divisible by q^m . Then we have

$$f(A * B) = (A_m + \phi(\overline{A}, \overline{B}), \overline{A} * \overline{B}).$$

So that f yields an isomorphism

$$\mathbb{Z}_{q^{m+1}}[T, T^{-1}]/(T - 1 + q) \rightarrow E(\mathbb{Z}_q[T, T^{-1}]/(T - 1 + q), \mathbb{Z}_q, \phi). \quad \blacksquare$$

Theorem 3.3 *For any positive integer q and m , the quandle $E = \mathbb{Z}_q[T, T^{-1}]/(1 - T)^{m+1}$ is an abelian extension of $X = \mathbb{Z}_q[T, T^{-1}]/(1 - T)^m$ over $\mathbb{Z}_q$: $E = E(X, \mathbb{Z}_q, \phi)$, for some $\phi \in Z_Q^2(X; \mathbb{Z}_q)$.*

Proof. Represent elements of E by $A = A_m(1 - T)^m + \dots + A_1(1 - T) + A_0$, where $A_j \in \mathbb{Z}_q$, $j = 0, \dots, m$. Define $f : E \rightarrow \mathbb{Z}_q \times X$ by

$$f(A) = (A_m \pmod{(1 - T)}, \overline{A} \pmod{(1 - T)^{m-1}}),$$

where $\overline{A} = \sum_{j=0}^{m-1} A_j(1 - T)^j$. Then for $A, B \in E$, the quandle operation is computed by

$$\begin{aligned} A * B &= TA + (1 - T)B \\ &= [1 - (1 - T)](A_m(1 - T)^m + \dots + A_1(1 - T) + A_0) \\ &\quad + (1 - T)(B_m(1 - T)^m + \dots + B_1(1 - T) + B_0) \\ &= (A_m - A_{m-1} + B_{m-1})(1 - T)^m + (A_{m-1} - A_{m-2} + B_{m-2})(1 - T)^{m-1} \\ &\quad + \dots + (A_1 - A_0 + B_0)(1 - T) + A_0 \\ &= (A_m - A_{m-1} + B_{m-1})(1 - T)^m + \sum_{j=0}^{m-1} (A_j - A_{j-1} + B_{j-1})(1 - T)^j, \end{aligned}$$

where A_{-1}, B_{-1} are understood to be zeros in the last summation, and the coefficients are in $\mathbb{Z}_q$. Note that in $\mathbb{Z}_q[T, T^{-1}]/(1 - T)^m$, we have

$$\overline{A} * \overline{B} = \sum_{j=0}^{m-1} (A_j - A_{j-1} + B_{j-1})(1 - T)^j.$$

Hence we have

$$f(A * B) = (A_m - A_{m-1} + B_{m-1}, \sum_{j=0}^{m-1} (A_j - A_{j-1} + B_{j-1})(1 - T)^j) \in \mathbb{Z}_q \times X.$$

Then we have

$$f(A * B) = (A_m + \phi(\overline{A}, \overline{B}), \overline{A} * \overline{B}),$$

where $\phi(\overline{A}, \overline{B}) = B_{m-1} - A_{m-1}$. Hence f yields an isomorphism

$$\mathbb{Z}_q[T, T^{-1}]/(1 - T)^{m+1} \rightarrow E(\mathbb{Z}_q[T, T^{-1}]/(1 - T)^m, \mathbb{Z}_q, \phi).$$

The cocycle ϕ has a similar description to the one in Theorem 3.2. Let

$$s : \mathbb{Z}_q[T, T^{-1}]/(1 - T)^m \rightarrow \mathbb{Z}_q[T, T^{-1}]/(1 - T)^{m+1}$$

be a set-theoretic section defined by

$$s \left(\sum_{j=0}^{m-1} A_j(1 - T)^j \pmod{(1 - T)^m} \right) = \sum_{j=0}^{m-1} A_j(1 - T)^j \pmod{(1 - T)^{m+1}}.$$

Then we have $\overline{s(X) * s(Y)} = \overline{s(X * Y)}$ for any $X, Y \in \mathbb{Z}_q[T, T^{-1}]/(1 - T)^m$. So that $[s(X) * s(Y) - s(X * Y)]$ is divisible by $(1 - T)^m$, and we have

$$\phi(\overline{A}, \overline{B}) = [s(X) * s(Y) - s(X * Y)]/(1 - T)^m \in \mathbb{Z}_q. \quad \blacksquare$$

Example 3.4 1. Consider the case $q = 2, m = 2$ in Theorem 3.2. In this case

$$\mathbb{Z}_4[T, T^{-1}]/(T + 1) = R_4, \quad \text{and}$$

$$\mathbb{Z}_8[T, T^{-1}]/(T + 1) = R_8 = E(R_4, \mathbb{Z}_2, \phi)$$

for some $\phi \in Z_Q^2(R_4; \mathbb{Z}_2)$. We obtain an explicit formula for this cocycle ϕ by computation:

$$\phi = \chi_{0,2} + \chi_{0,3} + \chi_{1,0} + \chi_{1,3} + \chi_{2,0} + \chi_{2,3} + \chi_{3,0} + \chi_{3,1},$$

where

$$\chi_{a,b}(x, y) = \begin{cases} 1 & \text{if } (x, y) = (a, b), \\ 0 & \text{if } (x, y) \neq (a, b) \end{cases}$$

denotes the characteristic function.

2. In case $m = 1$ and $q = 3$, the cocycle constructed is of the form

$$\phi = \chi_{0,1} + \chi_{1,2} + \chi_{2,0} + 2(\chi_{0,2} + \chi_{1,0} + \chi_{2,1}).$$

3. In case $m = 2$ and $q = 3$, the cocycle is

$$\begin{aligned} \phi &= \chi_{0,3} + \chi_{0,4} + \chi_{0,5} + 2\chi_{0,6} + 2\chi_{0,7} + 2\chi_{0,8} \\ &+ 2\chi_{1,0} + \chi_{1,4} + \chi_{1,5} + \chi_{1,6} + 2\chi_{1,7} + 2\chi_{1,8} \\ &+ 2\chi_{2,0} + 2\chi_{2,1} + \chi_{2,5} + \chi_{2,6} + \chi_{2,7} + 2\chi_{2,8} \\ &+ 2\chi_{3,0} + 2\chi_{3,1} + \chi_{3,5} + \chi_{3,6} + \chi_{3,7} + 2\chi_{3,8} \\ &+ 2\chi_{4,0} + 2\chi_{4,1} + 2\chi_{4,2} + \chi_{4,6} + \chi_{4,7} + \chi_{4,8} \\ &+ \chi_{5,0} + 2\chi_{5,1} + 2\chi_{5,2} + 2\chi_{5,3} + \chi_{5,7} + \chi_{5,8} \\ &+ \chi_{6,0} + 2\chi_{6,1} + 2\chi_{6,2} + 2\chi_{6,3} + \chi_{6,7} + \chi_{6,8} \\ &+ \chi_{7,0} + \chi_{7,1} + 2\chi_{7,2} + 2\chi_{7,3} + 2\chi_{7,4} + \chi_{7,8} \\ &+ \chi_{8,0} + \chi_{8,1} + \chi_{8,2} + 2\chi_{8,3} + 2\chi_{8,4} + 2\chi_{8,5} \end{aligned}$$

4. Consider the case $q = 2$ and $m = 2$ in Theorem 3.3. The quandle $\mathbb{Z}_2[T, T^{-1}]/(1 - T)^2$ is isomorphic to R_4 by the correspondence $0 \leftrightarrow 0(1 - T) + 0, 1 \leftrightarrow 0(1 - T) + 1, 2 \leftrightarrow 1(1 - T) + 0$, and $3 \leftrightarrow 1(1 - T) + 1$. This is a special case of the isomorphism

$$\mathbb{Z}_n[T, T^{-1}]/(1 - T)^2 \cong \mathbb{Z}_{n^2}[T, T^{-1}]/(T - (kn + 1)) \quad \text{if } (n, k) = 1$$

given in [19]. Then the quandle $\mathbb{Z}_2[T, T^{-1}]/(1 - T)^3$ is an abelian extension $E(R_4; \mathbb{Z}_2, \phi')$ for some $\phi' \in Z_Q^2(R_4; \mathbb{Z}_2)$. Then the cocycle $\phi'(\overline{A}, \overline{B}) = B_1 - A_1$ is 1 if and only if the pair $(\overline{A}, \overline{B})$ has distinct coefficients for $(1 - T)$, and we obtain

$$\phi' = \chi_{0,2} + \chi_{2,0} + \chi_{1,2} + \chi_{2,1} + \chi_{0,3} + \chi_{3,0} + \chi_{1,3} + \chi_{3,1}.$$

The cocycles $\phi_0 = \chi_{2,1} + \chi_{2,3}$, $\phi_1 = \chi_{1,0} + \chi_{1,2}$, and the above ϕ are linearly independent (evaluate on the cycles defined in Remark 3.9 below), and $\phi' = \phi + \phi_0 + \phi_1$.

We recall from [7] that two surjective homomorphisms of quandles $\pi_j : E_j \rightarrow X$, $j = 1, 2$, are called *equivalent* if there is a quandle isomorphism $f : E_1 \rightarrow E_2$ such that $\pi_1 = \pi_2 f$. In particular, abelian extensions define surjective homomorphisms $E(X, A, \phi) = A \times X \rightarrow X$ defined by the projection onto the second factor. It was proved in [7] that two abelian extensions $E(X, A, \phi)$ and $E(X, A, \phi')$ are equivalent in the above sense if and only if ϕ is cohomologous to ϕ' .

Proposition 3.5 *The abelian extensions*

$$\mathbb{Z}_{q^{m+1}}[T, T^{-1}]/(T - 1 + q) = E(\mathbb{Z}_{q^m}[T, T^{-1}]/(T - 1 + q), \mathbb{Z}_q, \phi),$$

$$\mathbb{Z}_q[T, T^{-1}]/(1 - T)^{m+1} = E(\mathbb{Z}_q[T, T^{-1}]/(1 - T)^m, \mathbb{Z}_q, \phi')$$

are not trivial, i.e., not product quandles.

Proof. Direct computations show that the chains

$$c = (0, 1) + (q, q^{m-1} + q - 1) \in Z_2^Q(X; \mathbb{Z}_q) \quad \text{and}$$

$$c' = (0, 1) + (1 - T, (1 - T)^{m-1} + (1 - T) - 1) \in Z_2^Q(X; \mathbb{Z}_q)$$

are cycles for $X = \mathbb{Z}_{q^m}[T, T^{-1}]/(T - 1 + q)$ and $X = \mathbb{Z}_q[T, T^{-1}]/(1 - T)^m$, respectively.

Then it is computed that $\phi(c) = 1$ and $\phi'(c') = 1$, and hence ϕ and ϕ' are not coboundaries, and the result follows. ■

Remark 3.6 We remark here on consequences on dihedral quandles we derive from the above results. The quandle structure of a dihedral quandle R_n is defined using ring structure of $\mathbb{Z}_n$. The product quandle $R_m \times R_n$ is defined by component-wise operation, so that it is defined from the ring structure of $\mathbb{Z}_m \times \mathbb{Z}_n$ as well. Consequently, two quandles $R_m \times R_n$ and R_{mn} are isomorphic if $\mathbb{Z}_m \times \mathbb{Z}_n$ and $\mathbb{Z}_{mn}$ are isomorphic as rings. Hence if $n = p_1^{e_1} \dots p_k^{e_k}$ is the prime decomposition, then R_n is isomorphic to $R_{p_1^{e_1}} \times \dots \times R_{p_k^{e_k}}$. For $p = 2$, the result of this section shows that R_{p^e} is described successively as an extension of $R_{p^{e-1}}$.

The following lemma follows from definitions.

Lemma 3.7 *Let X, Y be quandles and A be an abelian group. If E is an abelian extension of X for $\phi \in Z_Q^2(X; A)$: $E = E(X, A, \phi)$, then $E \times Y$ is an abelian extension of $X \times Y$ for $p^\# \phi \in Z_Q^2(X \times Y; A)$: $E \times Y = E(X \times Y, A, p^\# \phi)$, where $p : X \times Y \rightarrow X$ is the projection to the first factor.*

Corollary 3.8 *For any positive integer n , $E = R_{4n}$ is an abelian extension $E = E(R_{2n}, \mathbb{Z}_2, \phi)$ of $X = R_{2n}$ for some cocycle $\phi \in Z_Q^2(R_{2n}; \mathbb{Z}_2)$.*

Proof. Let $2n = 2^m k$ for an odd integer k . Then $R_{2n} \cong R_{2^m} \times R_k$ by Remark 3.6, and by Lemma 3.7, $R_{4n} \cong R_{2^{m+1}} \times R_k$ is an abelian extension of R_{2n} if $R_{2^{m+1}}$ is an abelian extension of R_{2^m} . This follows from Theorem 3.2 since $R_{2^m} \cong \mathbb{Z}_{2^m}[T, T^{-1}]/(T + 1)$. ■

Remark 3.9 By Lemma 3.1 and Cor. 3.8, there is a cocycle $\phi \in Z_{\mathbb{Q}}^2(R_{4n}; \mathbb{Z}_2)$ such that R_{8n} is isomorphic to $E(R_{4n}, \mathbb{Z}_2, \phi)$.

Let $\phi_{0,1}, \phi_{1,0} \in Z_{\mathbb{Q}}^2(R_{4n}; \mathbb{Z}_2)$ be cocycles defined by

$$\phi_{0,1} = p^{\#}(\chi_{0,1} + \chi_{0,3}), \quad \text{and} \quad \phi_{1,0} = p^{\#}(\chi_{1,0} + \chi_{1,2}),$$

respectively, where $p : R_{4n} \rightarrow R_4$ is a natural map $p(x \bmod (4n)) = x \bmod (4)$. Here, it is known [3] that

$$\chi_{0,1} + \chi_{0,3}, \quad \text{and} \quad \chi_{1,0} + \chi_{1,2}$$

are cocycles in $Z_{\mathbb{Q}}^2(R_4; \mathbb{Z}_2)$. It is directly computed that

$$c_{0,1} = (0, 1) + (2, 1), \quad c_{1,0} = (1, 0) + (4n - 1, 0), \quad c'_{0,1} = (0, 1) + (2, 2n + 1) \in Z_2^{\mathbb{Q}}(R_{4n}; \mathbb{Z}_2)$$

are cycles. Then we have

$$\begin{aligned} \phi_{0,1}(c_{0,1}) &= 1, & \phi_{0,1}(c_{1,0}) &= 0, & \phi_{0,1}(c'_{0,1}) &= 1, \\ \phi_{1,0}(c_{0,1}) &= 0, & \phi_{1,0}(c_{1,0}) &= 1, & \phi_{1,0}(c'_{0,1}) &= 0, \\ \phi(c_{0,1}) &= 0, & \phi(c_{1,0}) &= 0, & \phi(c'_{0,1}) &= 1, \end{aligned}$$

Hence we see that the cocycles $\phi_{0,1}, \phi_{1,0}$, and ϕ are linearly independent.

In [21, 19], ranks of homology groups are determined for certain families of quandles.

4 Cocycle knot invariants as obstructions to extending colorings

Let K be a knot, and denote by $\Phi_K(X, \phi)$ the state-sum invariant of K with respect to a quandle X and a cocycle $\phi \in Z_{\mathbb{Q}}^2(X; A)$, where A is an abelian group. Let $E = E(X, A, \phi)$ be the abelian extension of X by ϕ . We characterize when the state-sum invariant defined from cocycles is non-trivial, if the cocycles used are those defined from abelian extensions. For characterizations on the triviality of colorings, see [16].

Let $\mathcal{C}$ be a coloring of K by X . Let E be a quandle with a surjective homomorphism $p : E \rightarrow X$. In this case E is called an *extension* of X . If there is a coloring $\mathcal{C}'$ of K by E such that for every arc a of K , it holds that $p(\mathcal{C}'(a)) = \mathcal{C}(a)$, then $\mathcal{C}'$ is called an *extension* of $\mathcal{C}$.

Theorem 4.1 *Let $C_0(K, X)$ be the constant term (a positive integer) of $\Phi_K(X, \phi)$, and $C(K, X)$ be the number of all colorings of K by X . Then the number of colorings of K by X that extend to colorings of K by $E(X, A, \phi)$ is equal to $C_0(K, X)$, and the number of colorings that do not extend is $C(K, X) - C_0(K, X)$.*

Proof. Let $\mathcal{C}$ be a coloring whose contribution to $\Phi_K(X, \phi)$ is 1. Fix this coloring in what follows. Pick a base point b_0 on a knot diagram of K . Let $x \in X$ be the color on the arc α_0 containing b_0 . Let $\alpha_i, i = 1, \dots, n$, be the set of arcs that appear in this order when the diagram K is traced in the given orientation of K , starting from b_0 . Pick an element $a \in A$ and give a color (a, x) on α_0 , so that we define a coloring $\mathcal{C}'$ by E on α_0 by $\mathcal{C}'(\alpha_0) = (a, x) \in E$. We try to extend it to the entire diagram by traveling the diagram from b_0 along the arcs $\alpha_i, i = 1, \dots, n$, in this order, by induction.

Suppose $\mathcal{C}'(\alpha_i)$ is defined for $0 \leq i < k$. Define $\mathcal{C}'(\alpha_{k+1})$ as follows. Suppose that the crossing τ_k separating α_k and α_{k+1} is positive, and the over-arc at τ_k is α_j . Let $\mathcal{C}'(\alpha_k) = (a, x)$ and $\mathcal{C}(\alpha_j) = y \in X$. Then we have $\mathcal{C}(\alpha_{k+1}) = x * y \in X$. Define $\mathcal{C}'(\alpha_{k+1}) = (a\phi(x, y), x * y)$ in this case.

Suppose that the crossing τ_k is negative. Let $\mathcal{C}'(\alpha_k) = (a, x)$ and $\mathcal{C}(\alpha_j) = y \in X$. Then if $\mathcal{C}(\alpha_{k+1}) = z$, then we have $z * y = x$. Define $\mathcal{C}'(\alpha_{k+1}) = (a\phi(z, y)^{-1}, z)$ in this case.

Define $\mathcal{C}'(\alpha_i)$ inductively for all $i = 0, \dots, n$. Regard α_0 as α_{n+1} , and repeat the above construction at the last crossing τ_n to come back to α_0 . By the construction we have $\mathcal{C}'(\alpha_{n+1}) = (a \prod_{\tau} B(\tau, \mathcal{C}), \mathcal{C}(\alpha_0))$, where $\prod_{\tau} B(\tau, \mathcal{C})$ is the state-sum contribution (the product of Boltzmann weights over all crossings) of $\mathcal{C}$. This contribution is equal to 1 by the assumption that $\prod_{\tau} B(\tau, \mathcal{C}) = 1$, and we have a well-defined coloring $\mathcal{C}'$. Hence this color extends to $E(X, A, \phi)$.

Conversely, if a coloring $\mathcal{C}$ by X extends to a coloring by $E(X, A, \phi)$, then from the above argument, we have that $(a, x) = (a \prod_{\tau} B(\tau, \mathcal{C}), x)$, if (a, x) is the color on the base point b_0 . Hence $\prod_{\tau} B(\tau, \mathcal{C}) = 1$. ■

Thus the non-trivial value of $\Phi_K(X, \phi)$ is the obstruction to extending colorings of K by X to $E(X, A, \phi)$, in the following sense: there is a coloring $\mathcal{C}$ of K by X which does not extend to a coloring by $E(X, A, \phi)$, if and only if $\Phi_K(X, \phi)$ is not a positive integer.

Corollary 4.2 *For $A = \mathbb{Z}_2$, $\Phi_K(X, \phi) = a + bt$ (where t is the variable, the generator of $\mathbb{Z}_2$) is determined by the number of colorings with X and E : a is the number of colorings of X that extend to colorings by E , and b is the number of those that do not.*

Example 4.3 For $X = \mathbb{Z}_2[T, T^{-1}]/(T^2 + T + 1)$ we have a cocycle $\phi = \prod_{a, b \neq T} \chi_{(a, b)} \in Z_Q^2(X; \mathbb{Z}_2)$ (see [3]), let $E = E(X, \mathbb{Z}_2, \phi)$. Then $\Phi_K(S_4, \phi) = a + bt$ has the above characterization.

Specifically, in [4], it was computed that among knots in the table up to 9 crossings, the state-sum invariant with the above quandle and the cocycle takes the value $4 + 12t$ for the knots

$$3_1, 4_1, 7_2, 7_3, 8_1, 8_4, 8_{11}, 8_{13}, 9_1, 9_{12}, 9_{13}, 9_{14}, 9_{21}, 9_{23}, 9_{35}, 9_{37},$$

and the value $16 + 48t$ for $8_{18}, 9_{40}$. Hence for the knots in the former list, the number of colorings by X which extend to those by E is 4 (trivial colorings, by a single color), and those that do not extend is 12 (all non-trivial colorings do not extend). For the knots in the latter list, there are 16 colorings that extend, and 48 colorings that do not.

We generalize the invariants to links component-wise. Let $L = K_1 \cup \dots \cup K_n$ be a link diagram. The following generalization of the state-sum invariant, which follows from Reidemeister moves, is suggested by Rourke and Sanderson in personal communications.

Lemma 4.4 *Let UK_i , $i = 1, \dots, n$, denote the set of under-crossings on K_i . This is the set of crossings such that the under-arcs belong to the component K_i . Then the state-sum $\Phi_i(L) = \sum_{\mathcal{C}} \prod_{\tau \in UK_i} B(\tau, \mathcal{C})$ is an invariant of a link L for each $i = 1, \dots, n$.*

The vector $\vec{\Phi}(L) = (\Phi_i(L))_{i=1}^n$ of the state-sum invariants is called the *component-wise* (quandle) cocycle invariant of L . We observe here that the theorem in this section applies to component-wise cocycle invariants.




Figure 2: A colored Whitehead link

Theorem 4.5 Let $\vec{\Phi}(L) = (\Phi_i(L))_{i=1}^n$ be the component-wise cocycle invariant of a link $L = K_1 \cup \dots \cup K_n$ with a quandle X and a cocycle $\phi \in Z_Q^2(X; A)$ for an abelian group A . Then $\Phi_i(L)$ is not a positive integer for some i if and only if there is a coloring of L by X that does not extend to a coloring of L by $E(X, A, \phi)$.

Example 4.6 In Fig. 2, a Whitehead link $L = K_1 \cup K_2$ is depicted. Let $\phi \in Z_Q^2(R_8; \mathbb{Z}_2)$ be the cocycle defined in Cor. 3.8, that is, ϕ defines the extension $E = R_{16} = E(R_8, \mathbb{Z}_2, \phi)$. We evaluate the component-wise cocycle invariant $\vec{\Phi}(L) = (\Phi_1(L), \Phi_2(L))$. Denote the generator of the coefficient group $\mathbb{Z}_2$ by t , so that $\mathbb{Z}_2 = \{1, t\}$ and the invariant takes the form of $\vec{\Phi}(L) = (A_1 + B_1t, A_2 + B_2t)$, where A_i, B_i ($i = 1, 2$) are positive integers.

The colors assigned to arcs are represented by the letters a through e . From the figure, it is seen that all the colors are determined by the colors a and b assigned to the top two arcs. It is seen by calculations that for any choice of two elements of R_8 for a and b , there is a unique coloring of L by R_8 that restricts to the chosen elements for a and b . Therefore, there are $8^2 = 64$ colorings of L by R_8 .

We show that the state-sum term $\prod_{\tau \in UK_1} B(\tau, \mathcal{C})$ is 1 if and only if a and b have the same parity (both even or both odd).

Suppose that a and b are both even, so that $a = 2\alpha$, $b = 2\beta$. Then one computes that $c = 4\beta - 2\alpha$, $d = 6\beta - 4\alpha$, and we obtain $e = 2\beta = b$. Similar computations show that $e = b$ if a and b are both odd. From the figure, the state-sum term for UK_1 is $\phi(a, b)\phi(a, e)^{-1}$, which is equal to $\phi(a, b)\phi(a, b)^{-1} = 1$, in this case. Suppose now that a and b have opposite parities. By setting $a = 2\alpha + 1$ and $b = 2\beta$ (and vice versa), we compute that $e = b + 4$, so that we obtain the state-sum term $\phi(a, b)\phi(a, e)^{-1} = \phi(a, b)\phi(a, b + 4)^{-1}$. We claim that this is t .

Using the formula at the end of the proof of Theorem 3.2, we have $\phi(a, b) = [s(a)*s(b) - s(a*b)]/8$ (additively). Here, $s(a)*s(b)$ is $2b - a$ computed modulo 16, and $s(a*b)$ is $2b - a$ computed modulo 8, then regarded as an element modulo 16. Since $a*(b + 4) = 2(b + 4) - a = (2b - a) + 8$ modulo

16, we have

$$\phi(a, b) - \phi(a, b + 4) = [s(a) * s(b) - s(a * b)]/8 - [s(a) * s(b) + 8 - s(a * b)]/8 = 1 \bmod (2)$$

written additively. This proves the above claim. There are 32 colorings with the same parity, and 32 with distinct parities. Hence we obtain $\vec{\Phi}(L) = (32 + 32t, 32 + 32t)$.

Theorem 4.5 implies that there are colorings by R_8 that do not extend to colorings by R_{16} . In fact, from the proof of Theorem 4.1, we see that 32 colorings having the same parity for a and b extend to R_{16} , and those 32 colorings with the opposite parities do not. This fact can be computed directly, and gives an alternate method of computing the above invariant using Cor. 4.2.

Acknowledgments JSC is being supported by NSF grant DMS-9988107. MS is being supported by NSF grant DMS-9988101.

References

- [1] Brieskorn, E., *Automorphic sets and singularities*, Contemporary math., 78 (1988), 45–115.
- [2] Brown, K. S., *Cohomology of groups*. Graduate Texts in Mathematics, 87. Springer-Verlag, New York-Berlin, 1982.
- [3] Carter, J.S.; Jelsovsky, D.; Kamada, S.; Langford, L.; Saito, M., *Quandle cohomology and state-sum invariants of knotted curves and surfaces*, preprint at <http://xxx.lanl.gov/abs/math.GT/9903135> .
- [4] Carter, J.S.; Jelsovsky, D.; Kamada, S.; Saito, M., *Computations of quandle cocycle invariants of knotted curves and surfaces*, Advances in math., 157 (2001) 36-94.
- [5] Carter, J.S.; Jelsovsky, D.; Kamada, S.; Saito, M., *Quandle homology groups, their betti numbers, and virtual knots*, J. of Pure and Applied Algebra, 157, (2001), 135-155.
- [6] Carter, J.S.; Kamada, S.; Saito, M., *Geometric interpretations of quandle homology*, J. of Knot Theory and its Ramifications, vol. 10, no. 3 (2001), 345-358.
- [7] Carter, J.S.; Kamada, S.; Saito, M., *Diagrammatic Computations for Quandles and Cocycle Knot Invariants*, preprint at <http://xxx.lanl.gov/abs/math.GT/0102092>.
- [8] Carter, J.S.; Saito, M., *Knotted surfaces and their diagrams*, the American Mathematical Society, 1998.
- [9] Fenn, R.; Rourke, C., *Racks and links in codimension two*, Journal of Knot Theory and Its Ramifications Vol. 1 No. 4 (1992), 343-406.
- [10] Fenn, R.; Rourke, C.; Sanderson, B., *Trunks and classifying spaces*, Appl. Categ. Structures 3 (1995), no. 4, 321–356.
- [11] Fenn, R.; Rourke, C.; Sanderson, B., *James bundles and applications*, preprint found at <http://www.maths.warwick.ac.uk/~bjs/> .

- [12] Flower, Jean, *Cyclic Bordism and Rack Spaces*, Ph.D. Dissertation, Warwick (1995).
- [13] Fox, R.H., *A quick trip through knot theory*, in *Topology of 3-Manifolds*, Ed. M.K. Fort Jr., Prentice-Hall (1962) 120–167.
- [14] Gerstenhaber, M.; Schack, S. D., *Bialgebra cohomology, deformations, and quantum groups*. Proc. Nat. Acad. Sci. U.S.A. 87 (1990), no. 1, 478–481.
- [15] Greene, M. T. *Some Results in Geometric Topology and Geometry*, Ph.D. Dissertation, Warwick (1997).
- [16] Inoue, A., *Quandle homomorphisms of knot quandles to Alexander quandles*, Preprint.
- [17] Joyce, D., *A classifying invariant of knots, the knot quandle*, J. Pure Appl. Alg., 23, 37–65.
- [18] Kauffman, L. H., *Knots and Physics*, World Scientific, Series on knots and everything, vol. 1, 1991.
- [19] Litherland, R.A.; Nelson, S., *The Betti numbers of some finite racks*, preprint at: <http://xxx.lanl.gov/abs/math.GT/0106165>.
- [20] Matveev, S., *Distributive groupoids in knot theory*, (Russian) Mat. Sb. (N.S.) 119(161) (1982), no. 1, 78–88, 160.
- [21] Mochizuki, T., *Some calculations of cohomology groups of finite Alexander quandles*, Preprint.
- [22] Rourke, C., and Sanderson, B., *There are two 2-twist-spun trefoils*, preprint at <http://xxx.lanl.gov/abs/math.GT/0006062> .